

PRIME MESSAGES

1. Immunization protects against several dangerous diseases. A child who is not immunised is more likely to become undernourished, disabled or die.
2. Immunization is urgent. All the primary immunizations should be completed in the first year of life.
3. Booster doses of immunization should be given as per schedule.
4. Prophylactic immunization should be given as and when required.
5. It is safe to immunize a child suffering from mild illness in consultation with a doctor.
6. Every woman between ages of 15 and 44 should be immunized against Tetanus and Rubella.

EARLY INTERVENTION SERIES



IMMUNIZATION



National Institute for the Empowerment of Persons with Intellectual Disabilities (Divyangjan)

Department of Empowerment of Persons with Disabilities (Divyangjan)
Ministry of Social Justice & Empowerment, Government of India
(An ISO 9001:2015 Institution)

Manovikas Nagar, Secunderabad 500 009. Telangana, INDIA

Phone: 040-27751741-745, Fax: 040-27750198

E-mail: dir-niepid@niepid.nic.in, website: www.niepid.nic.in

Prepared under project:
**Early Intervention to IUGR Children At Risk for
Developmental Delays**

1. Baby's Name: _____

2. Father's Name: _____

IMMUNIZATION RECORD

AGE	VACCINE	DUE ON	GIVEN ON
Birth	BCG		
	Oral Polio Vaccine- 1st dose		
	Hepatitis B Vaccine - 1 dose		
6 weeks	DPT - 1st dose		
	Oral Polio Vaccine - 2nd dose		
	Hepatitis B Vaccine - 2nd dose		
10 weeks	DPT - 2nd dose		
	Oral Polio Vaccine - 3rd dose		
14 weeks	DPT - 3rd dose		
	Oral Polio Vaccine - 4th dose		
6-9 months	Oral Polio Vaccine - 5th dose		
	Hepatitis B Vaccine - 3rd dose		
9 months	Measles Vaccine		
15-18 months	MMR (Measles, Mumps, Rubella)		
	DPT - 1st booster dose		
	Oral Polio Vaccine - 6th dose		
5 years	DPT - 2nd booster dose		
	Oral Polio Vaccine - 7th dose		
10 years	TT (Tetanus) - 3rd booster dose		
15-16 years	TT (Tetanus) - 4th booster dose		

IMMUNIZATION SCHEDULE

BIRTH	DCG, OPV	Oral Polio Vaccine (OPV)
6 Weeks	DPT, OPV	(DPT-Diphtheria, Pertussis, Tetanus)
10 weeks	DPT, OPV	
14 weeks	DTP, OPV	
9 months	Measles, OPV	(Recommended by WHO)
12-15 months	MMR	(No Booster Required)
15-18 months	DPT, OPV	(First Booster)
4-5 years	DPT, OPV	(Second Booster)
10 years	TT	
10 years	Typhoid	
16 years	TT	
16 years	Typhoid	