




MULTIPLE CHOICE TYPE QUESTIONS

For 2025 Exams - Mathematics (041) - Class 11

☑ Select the correct option in the followings. Each question carries 1 mark.

01. Domain of $f(x) = \frac{1}{x+1}$ is $x \in$
 (a) \mathbb{R} (Real nos.) (b) $\mathbb{R} - 1$ (c) $\mathbb{R} - \{-1\}$ (d) $(-\infty, -1)$
02. If $|x| - 4 \geq 0$, then $x \in$
 (a) $(-\infty, -4] \cup [4, \infty)$ (b) $[-4, 4]$ (c) $(-\infty, -4) \cup (4, \infty)$ (d) $(-4, 4)$
03. If $A = \{0, 1, 2\}$ and, $B = \{\alpha, \beta\}$, then no. of functions defined from A to B is
 (a) 64 (b) 9 (c) 8 (d) 6
04. For the function $y = |-x|$, we shall always have $y \in$
 (a) \mathbb{Z}^+ (b) $(-\infty, 0]$ (c) $[0, \infty)$ (d) \mathbb{Z}
05. If $A = \{0, 2, 4\}$, $B = \{1, 3\}$ then, a relation R defined from A to B, having maximum number of elements is given by
 (a) $B \times B$ (b) $A \times A$ (c) $A \times B$ (d) $B \times A$
06. For the sets $A = \{2, 3\}$ and $B = \{1, 5, 6\}$, the total number of relations from A to B will be
 (a) 64 (b) 16 (c) 8 (d) 9
07. Let $A = \{1, 2, 3\}$, $B = \{4, 5, 6\}$.
 For a relation $R' : B \rightarrow A$ defined as $R' = \{(x, y) : x \in B, y \in A; x \text{ is divisible by } y\}$, the roster form is given by
 (a) $\{(1, 4), (2, 4), (1, 5), (1, 6), (2, 6), (3, 6)\}$ (b) $\{(4, 1), (4, 2), (5, 1), (6, 1), (6, 2), (6, 3)\}$
 (c) $\{(4, 1), (4, 2), (5, 1), (6, 1), (6, 2)\}$ (d) $\{(4, 1), (4, 2), (6, 1), (6, 2), (6, 3)\}$
08. For the sets $A = \{2, 3\}$, $B = \{4, 5, 6\}$, the value of $n(A \times B)$ will be
 (a) 8 (b) 9 (c) 6 (d) 64
09. Domain of $f(x) = \frac{1}{[x]}$ is given by $x \in$
 (a) $x \in \mathbb{R}$ (b) $x \in \mathbb{Z}$ (c) $x \in \mathbb{R} - \mathbb{Z}$ (d) $x \in \mathbb{R} - [0, 1)$
10. Consider the graph shown.




Which function is represented by this graph?

11. $f(x) = \sqrt{[x] - x}$ is defined when $x \in$
 (a) \mathbb{R} (Real nos.) (b) \mathbb{Z} (c) $\mathbb{R} - \mathbb{R}^-$ (d) ϕ

12. For $f(x) = 3 - |x|$, the range is given by
 (a) $(-\infty, 3)$ (b) $[0, \infty)$ (c) $\mathbb{R} - \mathbb{R}^-$ (d) $(-\infty, 3]$
13. Domain of $f(x) = \sqrt{1+x^2}$ is
 (a) \mathbb{R} (Real nos.) (b) $[1, \infty)$ (c) $(-\infty, 1]$ (d) $(1, \infty)$
14. Domain of $f(x) = \frac{1}{\sqrt{x+|x|}}$ is
 (a) $x \in \mathbb{R}$ (b) $x \in \mathbb{R} - \mathbb{R}^+$ (c) $x \in \mathbb{R} - \{0\}$ (d) $x \in \mathbb{R}^+$
15. Let $A = \{1, 2, 3\}$, $B = \{4, 5, 6\}$. Let $S : A \rightarrow B$, where $S = \{(1, 4), (2, 5), (3, 6), (1, 5)\}$. Then, S is
 (a) a relation only (b) a function only
 (c) a relation and function both (d) neither relation nor function
16. For $R = \{(x, y) : y = |x - 1|, x \in \mathbb{Z} \text{ and } |x| < 3\}$, the domain of R is
 (a) $\{\pm 3, \pm 2, \pm 1\}$ (b) $\{0, 1, 2\}$ (c) $\{\pm 2, \pm 1, 0\}$ (d) $\{\pm 3, \pm 2, \pm 1, 0\}$
17. If $y = \sqrt{(9-x)^2}$, then $x \in$
 (a) \mathbb{R} (Real nos.) (b) $\mathbb{R} - \{9\}$ (c) $(9, \infty)$ (d) $[9, \infty)$
18. The number of relations defined on the set $A = \{x : x \in \mathbb{N}, x < 1\}$ is
 (a) 1 (b) 0 (c) 2 (d) 4
19. If $f(x) = \begin{cases} -1, & \text{if } x < 0 \\ 0, & \text{if } x = 0 \\ 1, & \text{if } x > 0 \end{cases}$, then domain of function $f(x)$ is
 (a) $\{-1, 0, 1\}$ (b) $\{-1, 1\}$ (c) $x \in \mathbb{Z}$ (integers) (d) $x \in \mathbb{R}$ (real nos.)
20. For $f(x) = \sqrt{(x+1)^2} + \sqrt{x^2 - 9}$, the domain is given by
 (a) $x \in \mathbb{R}$ (Real nos.) (b) $x \in (-\infty, 3)$ (c) $x \in [3, \infty)$ (d) $x \in (-\infty, -3] \cup [3, \infty)$
21. Domain of $f(x) = \frac{1}{\sqrt{x}}$ is
 (a) $x \in \mathbb{R}$ (Real nos.) (b) $x \in \mathbb{R} - \{0\}$ (c) $x \in [0, \infty)$ (d) $x \in (0, \infty)$
22. If $A = \{1, 2\}$, $B = \{5, 6, 7\}$ and $C = \{5, 6, 7, 8\}$, then which of the following is correct?
 (a) $n[A \times (B \cup C)] = 6$ (b) $n[A \times (B \cup C)] = 14$
 (c) $n[A \times (B \cap C)] = 8$ (d) $n[A \times (B \cup C)] = 8$
23. For the function $f(x) = -|5x - 3|$, the range is
 (a) $[0, \infty)$ (b) $(-\infty, 0)$ (c) $(-\infty, 0]$ (d) $(0, \infty)$
24. If $A \times B = \{(1, 2), (1, 4), (3, 2), (3, 4)\}$, then $B \times A =$
 (a) $\{(2, 1), (4, 1), (2, 3), (4, 3)\}$ (b) $\{(1, 2), (1, 4), (3, 2), (3, 4)\}$
 (c) $\{(1, 2), (3, 2)\}$ (d) $\{(1, 2), (1, 4), (3, 2)\}$
25. If $A \times A = \{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3)\}$, then $A =$
 (a) $\{1, 2, 3\}$ (b) $\{1, 2\}$ (c) $\{0, 1, 2, 3\}$ (d) $\{0, 1, 2\}$
26. If $(x - 1, 2x - y) = (-1, 3)$, then y^x is
 (a) -3 (b) 3 (c) 1 (d) 0
27. If R be a relation from a set A to itself, then
 (a) $R = A$ (b) $R = A^2$ (c) $R \subseteq A \times A$ (d) $R \subseteq A \times B$

28. If $2f(x) - 3f\left(\frac{1}{x}\right) = x^2$, ($x \neq 0$), then $f(1) =$
 (a) 1 (b) -1 (c) 0 (d) 2
29. Let $[.]$ be a greatest integer function. For $-\frac{\pi}{2} < x < \frac{\pi}{2}$, the range of the function $f(x) = [2x]$ is
 (a) $\{\pm 3, \pm 2, \pm 1, 0\}$ (b) $\{\pm 3, \pm 2, \pm 1\}$ (c) $\{-4, \pm 3, \pm 2, \pm 1, 0\}$ (d) $\{\pm 4, \pm 3, \pm 2, \pm 1, 0\}$
30. If $f(x) = \frac{x+3}{x-3}$ and $g(x) = \frac{3x+x^3}{1+3x^2}$, then $g(1) =$
 (a) $f(1)$ (b) $f(0)$ (c) $\{f(0)\}^2$ (d) $\{f(1)\}^2$
31. If $f(x) = \frac{|x|}{4+|x|}$, then domain of $f(x)$ is
 (a) $x \in (0, \infty)$ (b) $x \in (-\infty, 0)$ (c) $x \in [0, \infty)$ (d) $x \in (-\infty, \infty)$
32. Domain of $f(x) = \frac{1}{\sqrt{x-x^2}}$ is
 (a) $\mathbb{R} - [0, 1]$ (b) $\mathbb{R} - (0, 1)$ (c) $(0, 1)$ (d) $[0, 1]$
33. If $[x]^2 - 5[x] - 6 = 0$, where $[.]$ denote the greatest integer function, then $x \in$
 (a) $[-1, 0) \cup [6, 7]$ (b) $[-1, 0) \cup [6, 7)$ (c) $[-1, 0] \cup [6, 7)$ (d) $[-1, 6]$
34. If $A = \{4, 5\}$ and $B = \{1, 2, 3\}$, then
 (a) $(4, 1) \in A \times B$ (b) $(2, 5) \in A \times B$ (c) $(4, 1) \in B \times A$ (d) $(2, 9) \in B \times A$
35. The number of empty relations defined on an empty set is
 (a) 1 (b) 2 (c) 0 (d) 4
36. Let $f(x) = \sqrt{1+x^2}$, then
 (a) $f(xy) = f(x)f(y)$ (b) $f(xy) \geq f(x)f(y)$ (c) $f(xy) \leq f(x)f(y)$ (d) None of these
37. The domain and range of the real function f defined by $f(x) = \frac{4-x}{x-4}$ is given by
 (a) Domain = \mathbb{R} , Range = $\{-1, 1\}$ (b) Domain = $\mathbb{R} - \{1\}$, Range = \mathbb{R}
 (c) Domain = $\mathbb{R} - \{4\}$, Range = $\{-1\}$ (d) Domain = $\mathbb{R} - \{-4\}$, Range = $\{-1, 1\}$
38. Range of $y = \frac{3}{2-x^2}$ is
 (a) $x > \frac{3}{2}$ (b) $x \geq \frac{3}{2}$ (c) $y > \frac{3}{2}$ (d) $y \geq \frac{3}{2}$
39. If $f(x) = \frac{x-1}{x+1}$, then
 (a) $f\left(\frac{1}{x}\right) = f(x)$ (b) $f\left(\frac{1}{x}\right) = -f(x)$ (c) $f\left(\frac{1}{x}\right) = 2f(x)$ (d) $f\left(\frac{1}{x}\right) = -2f(x)$
40. Let f and g be two real functions defined by $f(x) = \sqrt{x-1}$ and $g(x) = -2x+3$.
 Then the domain of $\frac{g}{f}$ is
 (a) $(1, \infty)$ (b) $[1, \infty)$ (c) $\mathbb{R} - \left\{\frac{3}{2}\right\}$ (d) $(-\infty, 1]$
41. Consider the graph of the function $y = f(x)$ given below.




Then the domain and range of the function $y = f(x)$ will be


- (a) Domain : $[-1, 3] \cup \{4\}$; Range : $[-2, 8)$ (b) Domain : $[-2, 8)$; Range : $[-1, 3] \cup \{4\}$
 (c) Domain : $[-2, 8)$; Range : $[-1, 4]$ (d) Domain : $[-2, 4)$; Range : $[-1, 3] \cup \{8\}$

42. Let $f : \mathbb{R} \rightarrow [0, \infty)$ be a function defined by $f(x) = |x|$ and $g(x) = f(x+1) + f(x-1) \forall x \in \mathbb{R}$. Then the graph of $g(x)$ is


(a)




(b)



(c)



(d)



Question numbers 43 to 45 are Assertion and Reason based questions. Two statements are given, one labelled **Assertion (A)** and the other labelled **Reason (R)**. Select the correct answer from the codes (a), (b), (c) and (d) as given below.

- (a) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of Assertion (A).
 (b) Both Assertion (A) and Reason (R) are true and Reason (R) is **not** the correct explanation of Assertion (A).
 (c) Assertion (A) is true but Reason (R) is false.
 (d) Assertion (A) is false but Reason (R) is true.

43. **Assertion (A)** : If the functions defined by $f(x) = 3x^2 - 1$ and $g(x) = 3 + x$ are equal, then the set of values of x will be $\left\{\frac{4}{3}, -1\right\}$.

Reason (R) : The range of greatest integer function $f(x) = [x]$ is \mathbb{R} (set of real numbers).

44. **Assertion (A)** : The ordered pair $(5, 2)$ belongs to the relation $R = \{(x, y) : y = x - 5; x, y \in \mathbb{Z}\}$.

Reason (R) : If $A \times B = \{(a, x), (a, y), (b, x), (b, y)\}$, then $A = \{a, b\}$ and $B = \{x, y\}$.

45. **Assertion (A)** : Given that $(x - 2, y + 5)$ and $\left(-2, \frac{1}{3}\right)$ are two equal ordered pairs, then $2024x + 3y + 14 = 0$.

Reason (R) : If f and g are real functions defined by $f(x) = x^2 + 7$ and $g(x) = 3x + 5$, then $(f \cdot g)(x) = 3x^3 + 5x^2 + 21x + 35$.

CLASS XI BASED ON NCERT TEXTBOOK & LATEST CBSE SYLLABUS FOR 2024-25

Mathmission

UNLEASH THE POWER OF AI WITH QR CODES TO ACCESS MORE FREE RESOURCES

- COMPLETE THEORY WITH EXAMPLES
- SUBJECTIVE TYPE QUESTIONS
- COMPETENCY FOCUSED QUESTIONS
- MULTIPLE CHOICE QUESTIONS
- ASSERTION - REASON QUESTIONS
- CASE STUDY QUESTIONS
- PASSAGE - BASED QUESTIONS

O.P. GUPTA
INDIRA AWARD WINNER

GET BULK-ORDERS @ THE DISCOUNTED PRICES FOR SCHOOLS, TUITIONS & COACHING INSTITUTES

Buy now online

amazon

Flipkart

with Keen Support of THOUSANDS OF TEACHERS ACROSS THE GLOBE

MOST REPUTED REFRESHER BOOK OF MATHS (041)

Order your copy now @ Discounted Rates only on WhatsApp @ +91-9650350480

CLASS XI

FOR CBSE EXAMS (2024-25)

SOLUTIONS OF MATHMISSION

O.P. GUPTA
INDIRA AWARD WINNER

We have released Set of 2 Books for CBSE Class XI (Academic session 2024-25).

1. MATHMISSION FOR XI

- ☑ COMPLETE THEORY & EXAMPLES
- ☑ SUBJECTIVE TYPE QUESTIONS
- ☑ COMPETENCY FOCUSED QUESTIONS
 - ★ Multiple Choice Questions
 - ★ Assertion-Reason Questions
 - ★ Case-Study Questions
 - ★ Passage-Based Questions
- ☑ ANSWERS OF ALL QUESTIONS

2. SOLUTIONS OF MATHMISSION

- ☑ Step-by-step Detailed Solutions (For all Exercises of MATHMISSION)

This document contains MCQs for Mathematics (041) of class XI.

★ Answers / Solutions is available on YouTube channel – Mathematicia By O.P. Gupta
You can share this document with other students!

✍ With a lot of Blessings!

O.P. GUPTA

Author & Math Mentor
Indira Award Winner

📖 The O.P. Gupta Advanced Math Classes
@ **Think Academy**, Near Dhansa Bus Stand
Metro Station Gate No.3, Najafgarh, Delhi

📞 Telegram / WhatsApp : +919650350480

📺 YouTube.com/@theopgupta

Exclusive coaching for Maths (041)
By **O.P. GUPTA**

- ☑ CBSE XII
- ☑ CBSE XI
- ☑ CUET
- ☑ JEE - MAIN
- ☑ NDA

MATHEMATICIA BY O.P. GUPTA

...a name you can bank upon!



Feel Safe to **Share this Document** with other math scholars

CLICK NOW

TO Download



or, just type -
theopgupta.com

**FREE PDF TESTS AND
ASSIGNMENTS OF THE
CLASSES XII, XI & X**



To get **FREE PDF Materials**, join
WhatsApp Teachers Group
by Clicking on the Logo



If you are a **Student**, then you may join our **Students Group**

CLICK HERE FOR
**CLASSES
IX & X**

CLICK HERE FOR
**CLASSES
XI & XII**

You can add our WhatsApp no. **+919650350480** to your Groups also

Many Direct Questions from our Books have been asked in the recent CBSE Exams



Click on the
Book cover
to buy!

**MATHMISSION
FOR XII & XI**
2024-25 Edition

Buy our
books on
amazon
Flipkart

/theopgupta /theopgupta /theopgupta /@theopgupta

For Bulk Orders of our Books at Discounted Price, contact on +91-9650350480