


PREBOARD EXAMINATION-(2025-26)**CLASS: XII****SUBJECT: MATHEMATICS (041)****Time Allowed: 3 Hours****Maximum Marks: 80****General Instructions:**

1. This question paper contains five sections - A, B, C, D and E. Each section is compulsory. However, there are internal choices in some questions.
2. Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
3. Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
4. Section C has 6 Short Answer (SA)-type questions of 3 marks each.
5. Section D has 4 Long Answer (LA)-type questions of 5 marks each.
6. Section E has 3 source based/case based/passage based/integrated units of assessment (4 marks each) with sub parts.

Q.No.	<u>SECTION – A</u> (Multiple Choice Questions) Each question carries 1 mark	Marks
1.	If $\begin{bmatrix} 3p & -6 \\ 1 & 2 \end{bmatrix}$ is a singular matrix, then the value of 'p' is : (a)2 (b)3 (c)0 (d)-1	1
2.	If A is a square matrix of order 3 and $ -2A = k A $, the value of k is (a) -8 (b) 8 (c) 4 (d) -4	1
3.	The value of λ for which the projection of $\vec{a} = \lambda\hat{i} + \hat{j} + 4\hat{k}$ on $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$ is 4 units : (a) 4 (b) 5 (c) 3 (d) 6	1
4.	The function $f(x) = \begin{cases} \frac{\sin x}{x} + \cos x, & x \neq 0 \\ k, & x = 0 \end{cases}$ is continuous at $x=0$, then the value of k is a. 3 b. 2 c. 1 d. 1.5	1
5.	The derivative of $\sec(\tan^{-1} x)$ w.r.t x is : (a) $\frac{x}{1+x^2}$ (b) $\frac{1}{\sqrt{1+x^2}}$ (c) $\frac{x}{\sqrt{1+x^2}}$ (d) $x\sqrt{1+x^2}$	1

6.	<p>If 'm' and 'n' respectively are the degree and order of the differential equation $\frac{d}{dx} \left[\left(\frac{dy}{dx} \right)^3 \right] = 0$, then m+n is :</p> <p>(a) 1 (b) 2 (c) 3 (d) 4</p>	1
7.	<p>For an L.P.P the objective function is $Z = 4x + 3y$, and the feasible region determined by a set of constraints (linear inequalities) is shown in the graph</p> <p>Which one of the following is true :</p> <p>(a) Maximum value of Z is at R. (b) Maximum value of Z is at Q. (c) Value of Z at R is less than the value at P. (d) Value of Z at Q is less than the value at R.</p> 	1
8.	<p>The area of a parallelogram whose adjacent sides are $\hat{i} - 2\hat{j} + 3\hat{k}$ and $2\hat{i} + \hat{j} - 4\hat{k}$ is :</p> <p>(a) $5\sqrt{3}$ sq. units (b) $10\sqrt{3}$ sq. units (c) $5\sqrt{6}$ sq. units (d) $10\sqrt{6}$ sq. units</p>	1
9.	<p>The value of $\int_1^{\sqrt{3}} \frac{1}{1+x^2} dx$ is :</p> <p>(a) $\frac{\pi}{3}$ (b) $\frac{2\pi}{3}$ (c) $\frac{\pi}{6}$ (d) $\frac{\pi}{12}$</p>	1
10.	<p>Let A be a square matrix the AA^T is :</p> <p>(a) Singular matrix (b) symmetric matrix (c) skew-symmetric matrix (d) non-singular matrix</p>	1
11.	<p>The value of $\int \frac{(x+1)(x+\log x)^2}{x} dx$ is :</p> <p>(a) $\frac{(x+\log x)^3}{3} + C$ (b) $\frac{(x+\log x)^2}{2} + C$ (c) $\frac{1}{x+\log x} + C$ (d) $\frac{1}{(x+\log x)^2} + C$</p>	1
12.	<p>If $\begin{vmatrix} x+1 & x-1 \\ x-3 & x+2 \end{vmatrix} = \begin{vmatrix} 4 & -1 \\ 1 & 3 \end{vmatrix}$, then the value of x is :</p> <p>(a) 4 (b) 3 (c) 0 (d) 2</p>	1
13.	<p>If A is a square matrix of order 3×3 such that $adj A = 25$, then A is :</p>	1

	(a) 125 (b) 9 (c) 5 (d) $\frac{1}{5}$	
14.	If A and B are two independent events such that $P(A) = 0.4$, $P(B) = p$ and $P(A \cup B) = 0.6$, then the value of 'p' is : (a) $\frac{1}{2}$ (b) $\frac{1}{3}$ (c) $\frac{2}{3}$ (d) $\frac{1}{5}$	1
15.	The corner points of the feasible region determined by the system of linear constraints are (0,3), (1,1) and (3,0). Let $Z = px + qy$, where $p, q > 0$. Conditions on p and q so that the minimum of z occurs at (3,0) and (1,1). (a) $p = 3q$ (b) $2p = q$ (c) $p = 3q$ (d) $p = q$	1
16.	If $y = 1 - x + \frac{x^2}{2!} - \frac{x^3}{3!} + \frac{x^4}{4!} - \dots$ then $\frac{d^2y}{dx^2}$ is equal to (a) $-x$ (b) $-y$ (c) x (d) y	1
17.	If ' θ ' is the angle between two vectors $\vec{a}$ and $\vec{b}$, then $ \vec{a} \cdot \vec{b} = \vec{a} \times \vec{b} $ when ' θ ' is equal to (a) 0 (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{2}$ (d) π	1
18.	The direction ratios of the line $4x - 12 = 2y + 4 = 3z - 3$ are (a) 4,6,3 (b) 6,3,4 (c) 4,6,3 (d) 3,6,4	1

ASSERTION-REASON BASED QUESTIONS:

In the following questions, a statement of assertion (A) is followed by a statement of Reason (R).

Choose the correct answer out of the following choices.

- (a) Both A and R are true and R is the correct explanation of A.
- (b) Both A and R are true but R is not the correct explanation of A.
- (c) A is true but R is false.
- (d) A is false but R is true.

19.	<p>Assertion (A): The domain of the function $\sec^{-1}(2x+1)$ is $(-\infty, -1] \cup [0, \infty)$.</p> <p>Reason (R): $\sec^{-1}\left(-\frac{2}{\sqrt{3}}\right) = \frac{5\pi}{6}$</p>	1
-----	--	---

20.	<p>Assertion (A): The lines $\frac{x-5}{7} = \frac{y+2}{-5} = \frac{z}{1}$ and $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$ are perpendicular.</p> <p>Reason (R): Two lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$ and $\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$ are perpendicular if $a_1a_2 + b_1b_2 + c_1c_2 = 0$.</p>	1
-----	--	---

SECTION –B

(This section comprises of very short answer type questions (VSA) of 2 marks each)

21.	<p>Find the value of $\sin\left(\frac{\pi}{2} - \sin^{-1}\left(\frac{-\sqrt{3}}{2}\right)\right)$</p> <p style="text-align: center;">OR</p> <p>Show that the function $f: \mathbb{R} \rightarrow \mathbb{R}$, defined as $f(x) = x^2$, is neither one-one nor onto.</p>	2
22.	A particle moves along the curve $6y = x^3 + 2$. Find the points on the curve at which y-coordinate is changing 8 times as fast as the x-coordinate.	2
23.	<p>Evaluate : $\int \sin^3 x \, dx$</p> <p style="text-align: center;">OR</p> <p>Evaluate : $\int \frac{dx}{e^x + e^{-x}}$</p>	2
24.	If $y = 500e^{7x} + 600e^{-7x}$, then show that $\frac{d^2y}{dx^2} = 49y$.	2
25.	Find the maximum profit that a company can make, if the profit function is given by $P(x) = 72 + 42x - x^2$, where x is the number of units and P is profit in rupees.	2

SECTION –C

(This section comprises of very short answer type questions (SA) of 3 marks each)

26.	Evaluate: $\int \frac{(x-3)}{(x-1)^3} e^x dx$	3
27.	<p>A bag contains $(2n + 1)$ coins. It is known that $(n - 1)$ of these coins have a head on both sides, whereas the rest of the coins are fair. A coin is picked up at random from the bag and is tossed. If the probability that the toss results in a head is $\frac{31}{42}$, determine the value of n.</p> <p style="text-align: center;">OR</p> <p>An electric shop has two types of LED bulbs of equal quantity. The probability of an LED bulb lasting more than 6 months given that it is of type 1 is 0.7 and is given that it is</p>	3

	of type 2 is 0.4. Then find the probability that on LED bulb chosen uniformly at random lasts more than 6 months..	
28.	<p>Evaluate $\int_0^{\frac{\pi}{2}} \frac{\cos^2 x}{1+3 \sin^2 x} dx$</p> <p style="text-align: center;">OR</p> <p>Evaluate: $\int_0^{\pi} \frac{x \sin x}{1+\cos^2 x} dx$</p>	3
29.	<p>Find the particular solution of the differential equation $(1 + e^{2x}) dy + (1 + y^2)e^x dx = 0$, given that $y = 1$ when $x = 0$.</p> <p style="text-align: center;">OR</p> <p>Solve the differential equation $ye^{x/y} dx = (x e^{x/y} + y^2)dy$ ($y \neq 0$).</p>	3
30.	<p>Solve the following Linear Programming Problem graphically:</p> <p>Maximize. $Z=4x+y$</p> <p>Subject to $x + y \leq 50$; $3x + y \leq 90$; $x, y \geq 0$.</p>	3
31.	Evaluate: $\int \frac{x^2+x+1}{(x^2+1)(x+2)} dx$	3
<p><u>SECTION-D</u></p> <p>(This section comprises of long answer type questions (LA) of 5 marks each)</p>		
32.	Using integration, find the area of the region enclosed between the circle $x^2 + y^2 = 16$ and the lines $y = 1$ and $y = 2$ in the first quadrant.	5
33.	<p>Show that the relation R defined on the set $\mathbf{N} \times \mathbf{N}$ by $(a,b) R (c,d) \Rightarrow a^2 + d^2 = b^2 + c^2$ $\forall a, b, c, d \in \mathbf{N}$ is an equivalence relation</p> <p style="text-align: center;">OR</p> <p>Show that the relation R on the set $A = \{x \in \mathbf{Z} : 0 \leq x \leq 12\}$, given by $R = \{(a,b) : a-b \text{ is a multiple of } 4\}$ is an equivalence relation. Also write equivalence class[1].</p>	5
34.	<p>Use the product</p> $\begin{bmatrix} 1 & -1 & 2 \\ 0 & 2 & -3 \\ 3 & -2 & 4 \end{bmatrix} \begin{bmatrix} -2 & 0 & 1 \\ 9 & 2 & -3 \\ 6 & 1 & -2 \end{bmatrix}$ <p>to solve the system of equation</p> $x - y + 2z = 1; 2y - 3z = 1; 3x - 2y + 4z = 2$	5

35.	<p>Find the coordinates of the image of the point (2, 3, 4) with respect to the line $\vec{r} = (2\hat{j} + 4\hat{k}) + \gamma (2\hat{i} + 4\hat{j} + \hat{k})$; where γ is a scalar. Also, find the distance of the image from the origin.</p> <p style="text-align: center;">OR</p> <p>If with reference to the right handed system of mutually perpendicular unit vectors $\hat{i}, \hat{j}$ and $\hat{k}$, $\vec{\alpha} = 3\hat{i} - \hat{j}$, $\vec{\beta} = 2\hat{i} + \hat{j} - 3\hat{k}$, then express $\vec{\beta}$ in the form $\vec{\beta} = \vec{\beta}_1 + \vec{\beta}_2$, where $\vec{\beta}_1$ is parallel to $\vec{\alpha}$ and $\vec{\beta}_2$ is perpendicular to $\vec{\alpha}$.</p>	5
-----	--	---

SECTION-E

This section contains three Case-study / Passage based questions. First two questions have three sub-parts (i), (ii) and (iii) of marks 1, 1 and 2 respectively. Third question has two sub-parts of 2 marks each.

36.	<div style="text-align: center;">  </div> <p>A shopkeeper sells three types of flowers seeds A_1, A_2 and A_3. These are sold as mixture, where their proportions are 4:4:2 respectively. Also their germination rates are 45%, 60% and 35% respectively. Let A_1: seed A_1 is chosen, A_2: seed A_2 is chosen and A_3: seed A_3 is chosen.</p> <p>Also let E: seed germinates.</p> <p>(i) Find $P(A_1)$, $P(A_2)$ and $P(A_3)$.</p> <p>(ii) Write $P(E A_1) + P(E A_2) + P(E A_3)$.</p> <p>(iii) Calculate the probability of a randomly chosen seed to germinate. Express the answer in %.</p> <p style="text-align: center;">OR</p> <p>(iii) Calculate the probability that seed is of the type A_2 given that a randomly chosen seed does not germinate.</p>	4
-----	--	---

37.	<p>CASE STUDY II: Read the following passage and answer the questions given below.</p>  <p>A sports stadium is elliptical in shape. The district sports administration wants to design a rectangular football field with the maximum possible area. The football field is given by the graph of $\frac{x^2}{25} + \frac{y^2}{9} = 1$.</p> <p>(i) If the length and the breadth of the rectangular field be '2x' and '2y' respectively, then find the area function in terms of 'x'.</p> <p>(ii) Find the critical point of the function obtained in (i).</p> <p>(iii) Use first derivative test to find the length '2x' and width '2y' of the soccer field, that will maximize its area.</p> <p style="text-align: center;">OR</p> <p>(iii) Use second derivative test to find the length '2x' and width '2y' of the soccer field, that will maximize its area.</p>	4
38.	<p>CASE STUDY III : Read the following passage and answer the questions given below.</p> <p>An airplane is flying along the line $\vec{r} = \alpha (2\hat{i} + 3\hat{j} + 4\hat{k})$; where α is a scalar and another aeroplane is flying along the line $\vec{r} = (\hat{i} + \hat{j}) + \gamma (3\hat{j} + 2\hat{k})$; where γ is a scalar.</p> <p>(i) At what points on the line should they reach, so that distance between them is shortest</p> <p>(ii) Find the shortest possible distance between them.</p>	4
